

Non-Quadratic Gauge Fixing and Ghosts for Gauge Theories on the Hypersphere

F. T. Brandt^a and D. G. C. McKeon^{b,c}

^a *Instituto de Física, Universidade de São Paulo, São Paulo, SP 05315-970, Brazil*

^b *Department of Applied Mathematics, University of Western Ontario, London, ON N6A 5B7, Canada and*

^c *Department of Mathematics and Computer Science,
Algoma University, Sault St. Marie, ON P6A 2G4, Canada*

It has been suggested that using a gauge fixing Lagrangian that is not quadratic in a gauge fixing condition is most appropriate for gauge theories formulated on a hypersphere. We reexamine the appropriate ghost action that is to be associated with gauge fixing, applying a technique that has been used for ensuring that the propagator for a massless spin-two field is transverse and traceless. It is shown that this non-quadratic gauge fixing Lagrangian leads to two pair of complex Fermionic ghosts and two Bosonic real ghosts.

PACS numbers: 11.15.-q

It has been shown that the classical action for Yang-Mills theory on an n -dimensional hypersphere [1–5] is

$$S_{cl} = \int d\Omega \left[-\frac{1}{12} (F_{abc}^i)^2 \right], \quad (1)$$

where $d\Omega$ is a surface element of a hypersurface of unit radius in this n -dimensional Euclidean space with coordinates η_a ($a = 1, \dots, n$) and

$$F_{abc}^i = L_{ab}A_c^i + f^{ijk}\eta_a A_b^j A_c^k + (\text{cyc. perm.})$$

$$(L_{ab} = \eta_a \frac{\partial}{\partial \eta_b} - \eta_b \frac{\partial}{\partial \eta_a}) \quad (2)$$

This action is invariant under the two gauge transformations

$$\delta_\phi A_a^i = \eta_a \phi^i(\eta) \quad (3a)$$

$$\delta_\theta A_a^i = \eta_p L_{pa} \theta^i(\eta) + f^{ijk} A_a^j \theta^k(\eta). \quad (3b)$$

(These transformations are infinitesimal; this is adequate for our purpose.)

In the path-integral formulation of the generating functional, it has been pointed out [4, 5] that the most convenient gauge fixing Lagrangian is

$$\mathcal{L}_{gf} = \frac{1}{2\alpha} [(L_{ab} + \delta_{ab}) A_b^i] [(L_{ac} - (n-2)\delta_{ac}) A_c^i] \quad (4)$$

The technique used in Ref [5] to derive the ghost action associated with this gauge fixing does not seem to work for standard Yang-Mills theory in Euclidean space in the Lorenz-Feynman gauge (even though it possesses a BRST invariance [6]); consequently we are motivated to reexamine how the ghost contribution to the effective action arises when one uses the non-quadratic gauge fixing Lagrangian of Eq. (4). The approach of Refs. [7, 8] can be used. (In the two references the transverse-traceless propagator for a spin-two field and the propagator for a spontaneously broken gauge theory are considered using non-quadratic gauge fixing.)

The technique we use is a generalization of 't Hooft's approach [9] to derive the Faddeev-Popov ghost action [10]. We first insert the following constant factors

$$\det [(L_{ab} + \delta_{ab}) \eta_b \delta^{ij}] \det [(L_{ab} + \delta_{ab}) (\eta_p L_{pb} \delta^{ij} + f^{ipj} A_b^p)]$$

$$\times \int \mathcal{D}\phi_1 \mathcal{D}\theta_1 \delta \left[(L_{ab} + \delta_{ab}) \left(A_b^i + \eta_b \phi_1^i + \eta_p L_{pb} \theta_1^i + f^{ijk} A_b^j \theta_1^k \right) - p_a^i \right] \quad (5a)$$

and

$$\det [(L_{ab} - (n-2)\delta_{ab}) \eta_b \delta^{ij}] \det [(L_{ab} - (n-2)\delta_{ab}) (\eta_p L_{pb} \delta^{ij} + f^{ipj} A_b^p)]$$

$$\times \int \mathcal{D}\phi_2 \mathcal{D}\theta_2 \delta \left[(L_{ab} - (n-2)\delta_{ab}) \left(A_b^i + \eta_b \phi_2^i + \eta_p L_{pb} \theta_2^i + f^{ijk} A_b^j \theta_2^k \right) - q_a^i \right] \quad (5b)$$

into the path integral

$$Z[J_a^i] = \int \mathcal{D}A_a^i \exp \left\{ \int d\Omega \left[-\frac{1}{12} (F_{abc}^i)^2 + J_a^i A_a^i \right] \right\} \quad (6)$$

We follow this with an insertion of the constant factor

$$\int \mathcal{D}p_a^i \mathcal{D}q_a^i \exp \left(\frac{1}{2\alpha} \int d\Omega p_a^i g_a^i \right) \quad (7)$$

into $Z[J]$. If now we perform the gauge transformation

$$A_a^i \rightarrow A_a^i - \delta_{\phi_1} A_a^i - \delta_{\theta_1} A_a^i \quad (8)$$

in the resulting expression for $Z[J]$ and then do the integration over p_a^i and q_a^i , we end up with

$$\begin{aligned} Z[J_a^i] = & \int \mathcal{D}A_a^i \int \mathcal{D}\phi_1 \mathcal{D}\theta_1 \int \mathcal{D}\phi \mathcal{D}\theta \det [(L_{ab} + \delta_{ab}) \eta_b \delta^{ij}] \det [(L_{ab} - (n-2)\delta_{ab}) \eta_b \delta^{ij}] \\ & \times \det [(L_{ab} + \delta_{ab}) (\eta_p L_{pb} \delta^{ij} + f^{ipj} A_b^p)] \det [(L_{ab} - (n-2)\delta_{ab}) (\eta_p L_{pb} \delta^{ij} + f^{ipj} A_b^p)] \\ & \times \exp \int d\Omega \left[-\frac{1}{12} (F_{abc}^i)^2 + \frac{1}{2\alpha} [(L_{ab} + \delta_{ab}) A_b^i] \right. \\ & \times \left. [(L_{ac} - (n-2)\delta_{ac}) (A_c^i + \eta_c \phi^i + \eta_p L_{pc} \theta^i + f^{ijk} A_c^j \theta^k)] + J_a^i A_a^i \right]; \quad (\phi^i = \phi_2^i - \phi_1^i, \quad \theta^i = \theta_2^i - \theta_1^i). \quad (9) \end{aligned}$$

We have used the fact the the functional determinants in Eqs. (5a) and (5b) are gauge invariant.

Each of the functional determinants in Eq. (9) can be exponentiated using the relation

$$\int \mathcal{D}c_a \mathcal{D}\bar{c}_a \exp [\bar{c}_a M_{ab} c_b] = \det M, \quad (10)$$

where c_a and $\bar{c}_a$ are complex fermionic “ghost” fields. Consequently there are two pair of complex Fermionic “ghost” fields arising from the functional determinants appearing in Eq. (9); in addition there are the Bosonic “ghost” fields ϕ^i and θ^i . The functional integrals over ϕ_1 and θ_1 serve only to rescale the normalization function for $Z[J_a]$. The gauge fixing Lagrangian of Eq. (4) has thus served to complicate the propagator for A_a^i because of the nature of these terms that occur in the exponential appearing in Eq. (9) which are quadratic in the fields A_a^i ,

ϕ^i and θ^i . By a “completing the square” operation, these quadratic terms can be made diagonal in the fields A_a^i , θ^i and ϕ^i , thereby retaining the vector propagator of Refs. [4, 5], though at the expense of having more complicated interaction terms.

Acknowledgments

F. T. Brandt would like to thank CNPq for financial support and the hospitality of the Department of Applied Mathematics of the University of Western Ontario, Canada. D. G. C. McKeon would like to thank R. MacLeod for a helpful suggestion, the Instituto de Física da Universidade de São Paulo, for the hospitality, and FAPESP, Brazil.

-
- [1] P. A. M. Dirac, The Annals of Mathematics **36**, pp. 657 (1935), ISSN 0003486X;
 - [2] S. L. Adler, Phys. Rev. D **6**, 3445 (1972); D **8**, 2400 (1973).
 - [3] R. Jackiw and C. Rebbi, Phys. Rev. Lett. **37**, 172 (1976).
 - [4] I. T. Drummond and G. M. Shore, Annals of Physics **117**, 89 (1979), ISSN 0003-4916,
 - [5] G. M. Shore, Annals of Physics **117**, 121 (1979), ISSN 0003-4916,
 - [6] D. G. C. McKeon, Mod. Phys. Lett. A **6**, 2201 (1991).
 - [7] F. T. Brandt, J. Frenkel, and D. G. C. McKeon, Phys. Rev. **D76**, 105029 (2007), 0707.2590.
 - [8] F. T. Brandt and D. G. C. McKeon, Phys. Rev. **D79**, 087702 (2009), 0903.3173.
 - [9] G. 't Hooft, Nucl. Phys. **B33**, 173 (1971).
 - [10] L. D. Faddeev and V. N. Popov, Phys. Lett. **B25**, 29 (1967).